

Essentials of Statistics: Exercises

David Brink



Download free books at

bookboon.com

David Brink

Statistics – Exercises

Statistics – Exercises

© 2010 David Brink & Ventus Publishing ApS

ISBN 978-87-7681-409-0

Contents

1	Preface	5
2	Problems for Chapter 2: Basic concepts of probability theory	6
3	Problems for Chapter 3: Random variables	8
4	Problems for Chapter 4: Expected value and variance	9
5	Problems for Chapter 5: The Law of Large Numbers	10
6	Problems for Chapter 6: Descriptive statistics	11
7	Problems for Chapter 7: Statistical hypothesis testing	12
8	Problems for Chapter 8: The binomial distribution	13
9	Problems for Chapter 9: The Poisson distribution	14
10	Problems for Chapter 10: The geometrical distribution	15
11	Problems for Chapter 11: The hypergeometrical distribution	16
12	Problems for Chapter 12: The multinomial distribution	17
13	Problems for Chapter 13: The negative binomial distribution	18
14	Problems for Chapter 14: The exponential distribution	19
15	Problems for Chapter 15: The normal distribution	20
16	Problems for Chapter 16: Distributions connected to the normal distribution	21
17	Problems for Chapter 17: Tests in the normal distribution	22
18	Problems for Chapter 18: Analysis of variance (ANOVA)	24
19	Problems for Chapter 19: The chi-squared test	25
20	Problems for Chapter 20: Contingency tables	26
21	Problems for Chapter 21: Distribution-free tests	27
22	Solutions	29

1 Preface

This collection of *Problems with Solutions* is a companion to my book *Statistics*. All references here are to this compendium.

2 Problems for Chapter 2: Basic concepts of probability theory

Problem 1

A *poker hand* consists of five cards chosen randomly from an ordinary pack of 52 cards. How many different possible hands N are there?

Problem 2

What is the probability of having the poker hand *royal flush*, i.e. Ace, King, Queen, Jack, 10, all of the same suit?

Problem 3

What is the probability of having the poker hand *straight flush*, i.e. five cards in sequence, all of the same suit?

Problem 4

What is the probability of having the poker hand *four of a kind*, i.e. four cards of the same value (four aces, four 7s, etc.)?

Problem 5

What is the probability of having the poker hand *full house*, i.e. three of a kind plus two of a kind?

Problem 6

What is the probability of having the poker hand *flush*, i.e. five cards of the same suit?

Problem 7

What is the probability of having the poker hand *straight*, i.e. five cards in sequence?

Problem 8

What is the probability of having the poker hand *three of a kind*?

Problem 9

What is the probability of having the poker hand *two pair*?

Problem 10

What is the probability of having the poker hand *one pair*?

Problem 11

A red and a black die are thrown. What is the probability P of having at least ten? What is the conditional probability Q of having at least ten, given that the black die shows five? What is the conditional probability R of having at least ten, given that at least one of the dice shows five?


Problem 12

How many subsets with three elements are there of a set with ten elements? How many subsets

with seven elements are there of a set with ten elements?

Problem 13

In how many ways can a set with 30 elements be divided into three subsets with five, ten and fifteen elements, respectively?




www.sylvania.com

We do not reinvent
the wheel we reinvent
light.

Fascinating lighting offers an infinite spectrum of possibilities: Innovative technologies and new markets provide both opportunities and challenges. An environment in which your expertise is in high demand. Enjoy the supportive working atmosphere within our global group and benefit from international career paths. Implement sustainable ideas in close cooperation with other specialists and contribute to influencing our future. Come and join us in reinventing light every day.

Light is OSRAM

OSRAM
SYLVANIA 



3 Problems for Chapter 3: Random variables

Problem 14

Consider a random variable X with point probabilities $P(X = k) = 1/6$ for $k = 1, 2, 3, 4, 5, 6$. Draw the graph of X 's distribution function $F : \mathbb{R} \rightarrow \mathbb{R}$.

Problem 15

Consider a random variable Y with density function $f(x) = 1$ for x in the interval $[0, 1]$. Draw the graph of Y 's distribution function $F : \mathbb{R} \rightarrow \mathbb{R}$.

Problem 16

A red and a black die are thrown. Let the random variable X be the sum of the dice, and let the random variable Y be the difference (red minus black). Determine the point probabilities of X and Y . Are X and Y independent?

Problem 17

A continuous random variable X has density

$$f(x) = \begin{cases} e^{-x} & \text{for } x \geq 0 \\ 0 & \text{for } x < 0 \end{cases}$$

Determine the distribution function F . What is $P(X > 1)$?

4 Problems for Chapter 4: Expected value and variance

Problem 18

A red and a black die are thrown, and X denotes the sum of the two dice. What is X 's expected value, variance, and standard deviation? What fraction of the probability mass lies within one standard deviation of the expected value?

Problem 19

A red and a black die are thrown. Let the random variable X be the sum of the two dice, and let the random variable Y be the difference (red minus black). Calculate the covariance of X and Y . How does this agree with the result of Problem 16, where we showed that X and Y are independent?

5 Problems for Chapter 5: The Law of Large Numbers

Problem 20

Let X be a random variable with expected value μ and standard deviation σ . What does Chebyshev's Inequality say about the probability $P(|X - \mu| \geq n\sigma)$? For which n is Chebyshev's Inequality interesting?

Problem 21

A coin is tossed n times and the number k of heads is counted. Calculate for $n = 10, 25, 50, 100, 250, 500, 1000, 2500, 5000, 10000$ the probability P_n that k/n lies between 0.45 and 0.55. Determine if Chebyshev's Inequality is satisfied. What does the *Law of Large Numbers* say about P_n ? Approximate P_n by means of the *Central Limit Theorem*.

Problem 22

Let X be normally distributed with standard deviation σ . Determine $P(|X - \mu| \geq 2\sigma)$. Compare with Chebyshev's Inequality.

Problem 23

Let X be exponentially distributed with intensity λ . Determine the expected value μ , the standard deviation σ , and the probability $P(|X - \mu| \geq 2\sigma)$. Compare with Chebyshev's Inequality.

Problem 24

Let X be binomially distributed with parameters $n = 10$ and $p = 1/2$. Determine the expected value μ , the standard deviation σ , and the probability $P(|X - \mu| \geq 2\sigma)$. Compare with Chebyshev's Inequality.

Problem 25

Let X be Poisson distributed with intensity $\lambda = 10$. Determine the expected value μ , the standard deviation σ , and the probability $P(|X - \mu| \geq 2\sigma)$. Compare with Chebyshev's Inequality.

Problem 26

Let X be geometrically distributed with probability parameter $p = 1/2$. Determine the expected value μ , the standard deviation σ , and the probability $P(|X - \mu| \geq 2\sigma)$. Compare with Chebyshev's Inequality.

6 Problems for Chapter 6: Descriptive statistics

Problem 27

Ten observations x_i are given:

4, 7, 2, 9, 12, 2, 20, 10, 5, 9

Determine the median, upper, and lower quartile and the inter-quartile range.

Problem 28

Four observations x_i are given:

2, 5, 10, 11

Determine the mean, empirical variance, and empirical standard deviation.



Discover the truth at www.deloitte.ca/careers

Deloitte.

© Deloitte & Touche LLP and affiliated entities.



7 Problems for Chapter 7: Statistical hypothesis testing

Problem 29

In order to test whether a certain coin is fair, it is tossed ten times and the number k of heads is counted. Let p be the “head probability”. We wish to test the null hypothesis

$$\mathbf{H}_0 : p = \frac{1}{2} \text{ (the coin is fair)}$$

against the alternative hypothesis

$$\mathbf{H}_1 : p > \frac{1}{2} \text{ (the coin is biased)}$$

We fix a significance level of 5%. What is the significance probability P if the number of heads is $k = 8$? Which values of k lead to acceptance and rejection, respectively, of \mathbf{H}_0 ? What is the risk of an error of type I? What is the strength of the test and the risk of an error of type II if the true value of p is 0.75?

8 Problems for Chapter 8: The binomial distribution

Problem 30

What is the probability P_1 of having at least six heads when tossing a coin ten times?

Problem 31

What is the probability P_2 of having at least 60 heads when tossing a coin 100 times?

Problem 32

What is the probability P_3 of having at least 600 heads when tossing a coin 1000 times?

SIMPLY CLEVER

ŠKODA



We will turn your CV into an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand? We will appreciate and reward both your enthusiasm and talent. Send us your CV. You will be surprised where it can take you.

Send us your CV on www.employerforlife.com



9 Problems for Chapter 9: The Poisson distribution

Problem 33

In a certain shop, an average of ten customers enter per hour. What is the probability P that at most eight customers enter during a given hour?

Problem 34

What is the probability Q that at most 80 customers enter the shop from the previous problem during a day of 10 hours?

Problem 35

At the 2006 FIFA World Championship, a total of 64 games were played. The number of goals per game was distributed as follows:

8	games with	0	goals
13	games with	1	goal
18	games with	2	goals
11	games with	3	goals
10	games with	4	goals
2	games with	5	goals
2	games with	6	goals

Determine whether the number of goals per game may be assumed to be Poisson distributed.

10 Problems for Chapter 10: The geometrical distribution

Problem 36

A die is thrown until one gets a 6. Let V be the number of throws used. What is the expected value of V ? What is the variance of V ?

Problem 37

Assume W is geometrically distributed with probability parameter p . What is $P(W < n)$?

Problem 38

In order to test whether a given die is fair, it is thrown until a 6 appears, and the number n of throws is counted. How great should n be before we can reject the null hypothesis

H_0 : the die is fair

against the alternative hypothesis

H_1 : the probability of having a 6 is less than $1/6$

at significance level 5%?

I joined MITAS because
I wanted **real responsibility**

The Graduate Programme
for Engineers and Geoscientists
www.discovermitas.com



Real work
International opportunities
Three work placements



Month 16
I was a construction
supervisor in
the North Sea
advising and
helping foremen
solve problems





11 Problems for Chapter 11: The hypergeometrical distribution

Problem 39

At a lotto game, seven balls are drawn randomly from an urn containing 37 balls numbered from 0 to 36. Calculate the probability P of having exactly k balls with an even number for $k = 0, 1, \dots, 7$.

Problem 40

Determine the same probabilities as in the previous problem, this time using the normal approximation.

12 Problems for Chapter 12: The multinomial distribution

Problem 41

A die is thrown six times. What is the probability of having two 4s, two 5s, and two 6s?

Problem 42

A die is thrown six times. What is the probability of having all six different numbers of pips?

ie business school

#1 EUROPEAN BUSINESS SCHOOL
FINANCIAL TIMES 2013

#gobeyond

MASTER IN MANAGEMENT

Because achieving your dreams is your greatest challenge. IE Business School's Master in Management taught in English, Spanish or bilingually, trains young high performance professionals at the beginning of their career through an innovative and stimulating program that will help them reach their full potential.

- Choose your area of specialization.
- Customize your master through the different options offered.
- Global Immersion Weeks in locations such as London, Silicon Valley or Shanghai.

Because you change, we change with you.

www.ie.edu/master-management | mim.admissions@ie.edu |



13 Problems for Chapter 13: The negative binomial distribution

Problem 43

At the 2006 FIFA World Championship, a total of 64 games were played. The number of goals per game is given in Problem 35. Investigate whether the number of goals per game may be assumed to be negatively binomially distributed.

14 Problems for Chapter 14: The exponential distribution

Problem 44

A device contains two electrical components, A and B. The lifespans of A and B are both exponentially distributed with expected lifespans of five years and ten years, respectively. The device works as long as both components work. What is the expected lifespan of the device?

Problem 45

A device contains two electrical components, A and B. The lifespans of A and B are both exponentially distributed with expected lifespans of five years. The device works as long as at least one of the components works. What is the expected lifespan of the device?

15 Problems for Chapter 15: The normal distribution

Problem 46

Let X be a normally distributed random variable with expected value $\mu = 3$ and variance $\sigma^2 = 4$. What is $P(X \geq 6)$?

Problem 47

Let X be a normally distributed random variable with expected value $\mu = 5$. Assume $P(X \leq 0) = 10\%$. What is the variance of X ?

Problem 48

A normally distributed random variable X satisfies $P(X \leq 0) = 0.40$ and $P(X \geq 10) = 0.10$. What is the expected value μ and the standard deviation σ ?

Problem 49

Consider independent random variables $X \sim N(1, 3)$ and $Y \sim N(2, 4)$. What is $P(X + Y \leq 5)$?

16 Problems for Chapter 16: Distributions connected to the normal distribution

Problem 50

Let Q be χ^2 distributed with $df = 70$ degrees of freedom. What is the expected value and the variance? What is $P(Q < 100)$?

Problem 51

Let X_1, \dots, X_{10} be independent standard normally distributed random variables. What is $P(X_1^2 + \dots + X_{10}^2 \leq 15)$?

Problem 52

Let T be t distributed with $df = 4$ degrees of freedom. What is the expected value and the variance? What is $P(T < 2)$?

Problem 53

Let V be F distributed with five degrees of freedom in the numerator and seven degrees of freedom in the denominator. Determine x such that $P(V < x) = 90\%$.

no.1
nine years in a row

Sweden
Stockholm

STUDY AT A TOP RANKED INTERNATIONAL BUSINESS SCHOOL

Reach your full potential at the Stockholm School of Economics, in one of the most innovative cities in the world. The School is ranked by the Financial Times as the number one business school in the Nordic and Baltic countries.

Visit us at www.hhs.se

STOCKHOLM SCHOOL OF ECONOMICS



17 Problems for Chapter 17: Tests in the normal distribution

Problem 54

Suppose we have a sample x_1, \dots, x_{10} of 10 independent observations from a normal distribution with variance $\sigma^2 = 3$ and unknown expected value μ . Assume that the sample has mean $\bar{x} = 0.7$. Test the null hypothesis

$$H_0 : \mu = 0$$

Problem 55

How great should the size of the sample in the previous problem have been in order to be able to reject H_0 ?

Problem 56

Suppose we have a sample consisting of four observations

$$2, 5, 10, 11$$

from a normal distribution with unknown expected value μ and unknown variance σ^2 . Test the null hypothesis

$$H_0 : \mu = 0$$

against the alternative hypothesis

$$H_1 : \mu > 0$$

Problem 57

Suppose we have a sample consisting of four observations

$$2, 5, 10, 11$$

from a normal distribution with unknown expected value μ and unknown variance σ^2 . Test the null hypothesis

$$H_0 : \sigma^2 = 10$$

against the alternative hypothesis

$$H_1 : \sigma^2 > 10$$

Problem 58

Suppose we have a sample consisting of four observations

$$2, 5, 10, 11$$

from a normal distribution with unknown expected value μ_1 and unknown variance σ_1^2 . Moreover, let there be given a sample

$$8, 12, 15, 17$$

from another independent normal distribution with unknown expected value μ_2 and unknown variance σ_2^2 . The observations of the second sample are somewhat greater than the observations of the first sample, but the question is whether this difference is significant. Test the null hypothesis

$$\mathbf{H}_0 : \mu_1 = \mu_2$$

against the alternative hypothesis

$$\mathbf{H}_1 : \mu_1 < \mu_2$$

18 Problems for Chapter 18: Analysis of variance (ANOVA)

Problem 59

Consider three samples:

sample 1: 2, 5, 10, 11

sample 2: 7, 11, 14, 16

sample 3: 3, 8, 9, 12

It is assumed that the samples come from independent normal distributions with common variance. Let μ_i be the expected value of the i 'th normal distribution. Test the null hypothesis

$$H_0 : \mu_1 = \mu_2 = \mu_3$$

using analysis of variance (ANOVA).



#1
in eco-friendly
attitude

**STUDY AT
LINKÖPING UNIVERSITY, SWEDEN**
RANKED AMONG TOP 50 UNIVERSITIES UNDER 50

Interested in Strategy and Management in International Organisations? Kick-start your career with a master's degree from Linköping University, Sweden.

→ **Click here!**

 **Linköping University**

19 Problems for Chapter 19: The chi-squared test

Problem 60

In 1998, Danish newspaper subscriptions were distributed as follows (simplified):

	Share
Berlingske Tidende	14%
Politiken	17%
Jyllands-Posten	20%
Information	5%
B.T.	25%
Ekstra Bladet	19%

In a 2008 market analysis, 100 randomly chosen persons were asked about their subscriptions. The result was:

	Number
Berlingske Tidende	18
Politiken	13
Jyllands-Posten	22
Information	2
B.T.	16
Ekstra Bladet	29

Determine using a χ^2 -test if the shares of the various newspapers have changed significantly since 1998.

Problem 61

Continuing the previous problem, we wish to determine which newspaper shares have significantly increased or decreased.

Problem 62

Over one week in spring 2005, the number of cars crossing the bridge between Denmark and Sweden was counted. The result was:

	Number
Monday	13804
Tuesday	13930
Wednesday	13863
Thursday	14023
Friday	14345
Saturday	14944
Sunday	15044

Is there a significant difference between these numbers?

20 Problems for Chapter 20: Contingency tables

Problem 63

In an opinion poll, randomly chosen Danes and Swedes were asked about their opinion (“pro” or “contra”) about euthanasia. The result was:

	pro	contra
Danes	70	22
Swedes	85	46

Investigate whether there is a significant difference between Denmark and Sweden in opinions about euthanasia.

Problem 64

A new medicine is tested in an experiment involving 40 patients. During the experiment, the medicine is given to 20 randomly chosen patients, and the remaining 20 patients are given a placebo treatment. After the treatment, it is seen which patients are still ill. The result was:

	fit	ill
medicine	8	12
placebo	2	18

Investigate whether the medicine has had a significant positive effect.

Problem 65

In a sociological investigation, three men and three women are asked if they watch football regularly. All the men say yes, while all the women say no. Is this difference statistically significant?

21 Problems for Chapter 21: Distribution-free tests

Problem 66

In a biological experiment, ten plants are treated with a certain pesticide. Before the treatment, the numbers of plant lice x_i on each plant are counted. One week after the treatment, the numbers of plant lice y_i are counted again. The result was:

Plant no.	x_i	y_i
1	41	27
2	51	59
3	66	76
4	68	65
5	46	36
6	69	54
7	47	49
8	44	51
9	60	55
10	44	35

Use Wilcoxon’s distribution-free test to determine whether the pesticide had a significant (positive) effect.

“I studied English for 16 years but...
...I finally learned to speak it in just six lessons”
Jane, Chinese architect

ENGLISH OUT THERE

Click to hear me talking before and after my unique course download



Problem 67

The experiment of the previous problem is repeated, this time with 100 plants. The statistics now become

$$t_+ = 3045 \text{ and } t_- = 2005$$

Is there a significant effect now?

Problem 68

A car factory counts the number of defects in randomly chosen cars from two different production lines.

Car no.	Line 1	Line 2
1	170	79
2	197	126
3	50	189
4	151	137
5	94	167
6	46	188
7	173	54
8	26	155
9	118	82
10	171	218
11	70	242
12	146	
13	55	
14	132	

Use a distribution-free test to determine if there are significantly more defects on one of the two production lines.

Problem 69

The sample from the previous problem is enlarged such that there are now $n = 100$ observations from Line 1 and $m = 50$ observations from Line 2. The statistic is found to be

$$t_x = 7002$$

Is there now a significant difference between the numbers of defects on the two lines?

22 Solutions

Solution of Problem 1

As stated in section 2.5 of the *Compendium*, this number can be computed as a binomial coefficient:

$$N = \binom{52}{5} = 2598960$$

Solution of Problem 2

We know the number of possible poker hands N from Problem 1. Of these, only four hands are royal flush. Therefore, the probability becomes

$$P = \frac{4}{N} = \frac{4}{2598960} = \frac{1}{649740} \approx 0.00015\%$$

Solution of Problem 3

We know the number of possible poker hands N from Problem 1. We have to calculate the number n of hands with straight flush. There are four possibilities for the suit. There are ten possibilities for the value of the highest card (from 5 to ace). This gives

$$4 \cdot 10 = 40$$

possibilities. However, we have to subtract the number of hands with royal flush (Problem 2) from this number. In total we get

$$n = 40 - 4 = 36$$

hands with straight flush. The probability of a straight flush thus becomes

$$P = \frac{n}{N} = \frac{36}{2598960} \approx \frac{1}{72193} \approx 0.0014\%$$

Solution of Problem 4

We know the number of possible poker hands N from Problem 1. We have to calculate the number n of hands with four of a kind. There are 13 possible values (ace, king, queen, etc.) of the four cards, and moreover 48 possibilities for the fifth card. In total this is

$$n = 13 \cdot 48 = 624$$

hands with four of a kind. The probability of four of a kind thus becomes

$$P = \frac{n}{N} = \frac{624}{2598960} = \frac{1}{4165} \approx 0.024\%$$

Solution of Problem 5

We know the number of possible poker hands N from Problem 1. We have to calculate the number n of hands with “full house”. There are 13 possibilities for the value of the group of three cards.

Moreover, there are 12 possibilities for the value of the pair of cards. Finally, one has to take into account that if we have, say, three aces and two kings, then the three aces can be chosen in

$$\binom{4}{3} = 4$$

ways, and the two kings can be chosen in

$$\binom{4}{2} = 6$$

ways. In total there are

$$n = 13 \cdot 12 \cdot 4 \cdot 6 = 3744$$

hands with full house. The probability of full house thus becomes

$$P = \frac{n}{N} = \frac{3744}{2598960} \approx \frac{1}{694} \approx 0.14\%$$

Solution of Problem 6

We have to calculate the number n of hands with a flush. There are four possibilities for the suit (spades, hearts, diamonds, clubs). Out of 13 cards of the same suit, one can choose five cards in

$$\binom{13}{5} = 1287$$

Excellent Economics and Business programmes at:



university of
 groningen




“The perfect start of a successful, international career.”

CLICK HERE
 to discover why both socially and academically the University of Groningen is one of the best places for a student to be

www.rug.nl/feb/education



ways. This gives

$$4 \cdot 1287 = 5148$$

possibilities. However, we have to subtract the number of hands with a straight flush (Problem 3) and the number of hands with a royal flush (Problem 2) from this number. In total we get

$$n = 5148 - 36 - 4 = 5108$$

hands with a flush. The probability of a flush thus becomes

$$P = \frac{n}{N} = \frac{5108}{2598960} \approx \frac{1}{509} \approx 0.20\%$$

Solution of Problem 7

We determine the number n of hands with a straight. For the value of the highest card, there are ten possibilities (from 5 to ace). For each of the five cards, there are four possibilities for the suit (spades, hearts, diamonds, clubs). This gives

$$10 \cdot 4 \cdot 4 \cdot 4 \cdot 4 \cdot 4 = 10240$$

possibilities. However, we have to subtract the number of hands with a straight flush (Problem 3) and the number of hands with a royal flush (Problem 2) from this number. In total we get

$$n = 10240 - 36 - 4 = 10200$$

hands with a straight. The probability of a straight thus becomes

$$P = \frac{n}{N} = \frac{10200}{2598960} \approx \frac{1}{255} \approx 0.39\%$$

Solution of Problem 8

We have to calculate the number n of hands with three of a kind. There are 13 possibilities for the value of the three cards. If we have, say, three aces, then these aces can be chosen in

$$\binom{4}{3} = 4$$

different ways. Moreover, there are

$$\binom{12}{2} = 66$$

possibilities for the values of the remaining two cards, and four possibilities for the suit of each of these. In total there are

$$n = 13 \cdot 4 \cdot 66 \cdot 4 \cdot 4 = 54912$$

hands with three of a kind. The probability of three of a kind thus becomes

$$P = \frac{n}{N} = \frac{54912}{2598960} \approx \frac{1}{47} \approx 2.1\%$$

Solution of Problem 9

We have to calculate the number n of hands with two pair. There are

$$\binom{13}{2} = 78$$

possibilities for the values of the two pair. For both of these two values, a pair can be chosen in

$$\binom{4}{2} = 6$$

different ways. There are 44 possibilities for the fifth card. In total there are

$$n = 78 \cdot 6 \cdot 6 \cdot 44 = 123552$$

hands with two pair. The probability of two pair thus becomes

$$P = \frac{n}{N} = \frac{123552}{2598960} \approx \frac{1}{21} \approx 4.8\%$$

Solution of Problem 10

We have to calculate the number n of hands with one pair. There are 13 possibilities for the value of the pair. One pair of given value can be chosen in

$$\binom{4}{2} = 6$$

different ways. For the values of the remaining three cards there are

$$\binom{12}{3} = 220$$

possibilities (the remaining three cards must have values different from each other and different from the pair). Finally there are four possibilities for the suit of each of the remaining three cards. In total this gives

$$n = 13 \cdot 6 \cdot 220 \cdot 4 \cdot 4 \cdot 4 = 1098240$$

hands with “one pair”. The probability of one pair thus becomes

$$P = \frac{n}{N} = \frac{1098240}{2598960} \approx 42\%$$

Solution of Problem 11

There is a total of 36 different sample points (i, j) . Of these, there are six sample points with sum at least ten, namely $(4, 6)$, $(5, 5)$, $(5, 6)$, $(6, 4)$, $(6, 5)$ and $(6, 6)$. The probability thus becomes

$$P = \frac{6}{36} = \frac{1}{6} \approx 17\%$$

There are six possible sample points $(i, 5)$ where the black die shows 5. Of these, there are two sample points where the sum is at least ten, namely $(5, 5)$ and $(6, 5)$. The conditional probability thus becomes

$$Q = \frac{2}{6} = \frac{1}{3} \approx 33\%$$

There are 11 possible sample points where at least one of the dice shows 5. Of these, there are three sample points where the sum is at least ten, namely $(5, 5)$, $(5, 6)$ and $(6, 5)$. The conditional probability thus becomes

$$R = \frac{3}{11} \approx 27\%$$


Solution of Problem 12

The number of subsets with three elements equals the binomial coefficient

$$\binom{10}{3} = \frac{10!}{3!7!} = \frac{10 \cdot 9 \cdot 8}{3 \cdot 2 \cdot 1} = 120$$

The number of subsets with seven elements is the same:

$$\binom{10}{7} = \binom{10}{3} = 120$$



In the past four years we have drilled

89,000 km

That's more than **twice** around the world.

Who are we?
 We are the world's largest oilfield services company¹. Working globally—often in remote and challenging locations—we invent, design, engineer, and apply technology to help our customers find and produce oil and gas safely.

Who are we looking for?
 Every year, we need thousands of graduates to begin dynamic careers in the following domains:

- Engineering, Research and Operations
- Geoscience and Petrotechnical
- Commercial and Business

What will you be?

careers.slb.com

Schlumberger

¹Based on Fortune 500 ranking 2011. Copyright © 2015 Schlumberger. All rights reserved.



Solution of Problem 13

The answer is the multinomial coefficient

$$\binom{30}{5 \ 10 \ 15} = \frac{30!}{5!10!15!} = 465817912560$$

Solution of Problem 14

The distribution function F is a step function given by

$$F(x) = \begin{cases} 0 & \text{for } x < 1 \\ 1/6 & \text{for } 1 \leq x < 2 \\ 2/6 & \text{for } 2 \leq x < 3 \\ 3/6 & \text{for } 3 \leq x < 4 \\ 4/6 & \text{for } 4 \leq x < 5 \\ 5/6 & \text{for } 5 \leq x < 6 \\ 1 & \text{for } x \geq 6 \end{cases}$$

Solution of Problem 15

The distribution function F is the continuous function given by

$$F(x) = \begin{cases} 0 & \text{for } x < 0 \\ x & \text{for } 0 \leq x \leq 1 \\ 1 & \text{for } x > 1 \end{cases}$$

Solution of Problem 16

X takes values in the set $\{2, 3, 4, \dots, 12\}$. The point probabilities are

$$P(X = k) = \begin{cases} 1/36 & \text{for } k = 2 \\ 2/36 & \text{for } k = 3 \\ 3/36 & \text{for } k = 4 \\ 4/36 & \text{for } k = 5 \\ 5/36 & \text{for } k = 6 \\ 6/36 & \text{for } k = 7 \\ 5/36 & \text{for } k = 8 \\ 4/36 & \text{for } k = 9 \\ 3/36 & \text{for } k = 10 \\ 2/36 & \text{for } k = 11 \\ 1/36 & \text{for } k = 12 \end{cases}$$

Y takes values in the set $\{-5, -4, \dots, 4, 5\}$. The point probabilities are

$$P(Y = k) = \begin{cases} 1/36 & \text{for } k = -5 \\ 2/36 & \text{for } k = -4 \\ 3/36 & \text{for } k = -3 \\ 4/36 & \text{for } k = -2 \\ 5/36 & \text{for } k = -1 \\ 6/36 & \text{for } k = 0 \\ 5/36 & \text{for } k = 1 \\ 4/36 & \text{for } k = 2 \\ 3/36 & \text{for } k = 3 \\ 2/36 & \text{for } k = 4 \\ 1/36 & \text{for } k = 5 \end{cases}$$

If we know, say, that X takes the value 12, then we can conclude that Y takes the value 0 (since both dice in that case show 6). Consequently, X and Y are *not* independent.

Solution of Problem 17

The distribution function is found by integrating the density:

$$F(x) = \int_{-\infty}^x e^{-t} dt = \begin{cases} 0 & \text{for } x \leq 0 \\ 1 - e^{-x} & \text{for } x > 0 \end{cases}$$

Moreover:

$$P(X > 1) = 1 - F(1) = e^{-1}$$

Solution of Problem 18

We know the point probabilities from Problem 16. Since the point probabilities are symmetrical around 7, we get at once

$$E(X) = 7$$

Let us compute the variance using the formula $\text{var}(X) = E(X^2) - E(X)^2$ from section 4.5. We get

$$E(X^2) = \sum_{k=2}^{12} P(X = k) \cdot k^2 = \frac{1974}{36} = 54.8$$

and thus

$$\text{var}(X) = E(X^2) - E(X)^2 = 54.8 - 7^2 = 5.8$$

The standard deviation is the square root of the variance:

$$\sigma = \sqrt{\text{var}(X)} = \sqrt{5.8} = 2.4$$

Within 1 standard deviation around the the expected value, that is in the interval from $7 - 2.4$ to $7 + 2.4$, we have

$$\sum_{k=5}^9 P(X = k) = \frac{4}{36} + \frac{5}{36} + \frac{6}{36} + \frac{5}{36} + \frac{4}{36} = \frac{24}{36}$$

i.e. two thirds of the probability mass (as predicted in section 4.4).

Solution of Problem 19

Let us find the covariance using the formula

$$\text{Cov}(X, Y) = E(X \cdot Y) - E(X) \cdot E(Y)$$

from section 4.6. From the point probabilities it appears that

$$E(X) = 7 \quad \text{and} \quad E(Y) = 0$$

It is somewhat tedious to calculate $E(X \cdot Y)$. There is a total of 36 possible sample points (r, s) when the red and the black die are thrown. For any given sample point (r, s) , we know that X takes the value $r + s$, whereas Y takes the value $r - s$. The product $X \cdot Y$ thus takes the value $(r + s)(r - s) = r^2 - s^2$. The expected value of $X \cdot Y$ is the mean of these 36 values, i.e.

$$E(X \cdot Y) = \frac{1}{36} \sum_{r=1}^6 \sum_{s=1}^6 (r^2 - s^2) = 0$$

American online

LIGS University

is currently enrolling in the
Interactive Online **BBA, MBA, MSc,**
DBA and PhD programs:

- ▶ enroll **by September 30th, 2014** and
- ▶ **save up to 16%** on the tuition!
- ▶ pay in 10 installments / 2 years
- ▶ Interactive **Online education**
- ▶ visit www.ligsuniversity.com to find out more!

Note: LIGS University is not accredited by any nationally recognized accrediting agency listed by the US Secretary of Education. More info [here](#).





We therefore find the covariance

$$\text{Cov}(X, Y) = E(X \cdot Y) - E(X) \cdot E(Y) = 0 - 7 \cdot 0 = 0$$

The covariance of *independent* random variables is always 0 (section 4.6). We have shown in 16 that X and Y are *not* independent. This problem therefore shows that we *cannot* conclude conversely that the variables are independent when the covariance is 0.

Solution of Problem 20

Chebyshev’s Inequality says

$$P(|X - \mu| \geq n\sigma) \leq \frac{1}{n^2}$$

In words this means that the probability that X takes values more than n standard deviations away from its expected value is small when n is large. Of course this is only interesting for $n > 1$ since every probability a priori is at most 1.

Solution of Problem 21

The number k is obviously binomially distributed with parameters n and $p = 1/2$. It is seen that k/n lies in the interval $[0.45 ; 0.55]$ if and only if k lies in the interval $[0.45n ; 0.55n]$. We may therefore calculate P_n using the formula for the point probabilities of the binomial distribution. The results are seen here:

n	P_n
10	0.25
25	0.38
50	0.52
100	0.73
250	0.886
500	0.978
1000	0.9986
2500	0.99999949
5000	0.999999999987
10000	0.9999999999999999999987

Since k has expected value $np = n/2$ and standard deviation $\sqrt{npq} = \sqrt{n/4} = \sqrt{n}/2$, it follows that k/n will have expected value $\mu = 1/2$ and standard deviation $\sigma = 1/2\sqrt{n}$. Note that the standard deviation converges to 0 as n goes to infinity.

Chebyshev’s Inequality gives

$$P_n = P(|k/n - 1/2| \leq 0.05) \geq 1 - \frac{\sigma^2}{(0.05)^2} = 1 - \frac{100}{n}$$

which is only interesting for $n > 100$. Here a table shows the right-hand side of Chebyshev’s

Inequality:

n	$1 - 100/n$
10	–9
25	–3
50	–1
100	0
250	0.6
500	0.8
1000	0.9
2500	0.96
5000	0.98
10000	0.99

By comparison with the first table, it is seen that Chebyshev’s Inequality really is satisfied.

The Law of Large Numbers says that P_n converges to 100% as n goes to infinity. We note that this is a direct consequence of Chebyshev’s Inequality and also appears very clearly from both of the above tables.

The Central Limit Theorem tells that the distribution of k/n approaches a normal distribution when n goes to infinity. Thus P_n can be approximated by

$$Q_n := \Phi\left(\frac{\sqrt{n}}{10}\right) - \Phi\left(-\frac{\sqrt{n}}{10}\right) = 1 - 2 \cdot \Phi\left(-\frac{\sqrt{n}}{10}\right)$$

since 0.05 equals $\sqrt{n}/10$ times the standard deviation σ . Here a table shows how well Q_n approximates P_n :

n	Q_n
10	0.25
25	0.38
50	0.52
100	0.68
250	0.886
500	0.975
1000	0.9984
2500	0.99999943
5000	0.999999999985
10000	0.999999999999999999999985

Solution of Problem 22

The probability is

$$P(|X - \mu| \geq 2\sigma) = 2 \cdot \Phi(-2) = 2.6\%$$

In contrast, Chebyshev’s Inequality only gives the weaker statement

$$P(|X - \mu| \geq 2\sigma) \leq \frac{\sigma^2}{(2\sigma)^2} = 25\%$$

Solution of Problem 23

The expected value of X is $1/\lambda$ and the standard deviation is $\sigma = \sqrt{1/\lambda^2} = 1/\lambda$. The probability that X takes a value more than two standard deviations from μ is

$$P(X \geq 3/\lambda) = 1 - F(3/\lambda) = \exp(-3) = 5.0\%$$

where we have used the distribution function of the exponential distribution, $F(x) = 1 - \exp(-\lambda x)$. In contrast, Chebyshev's Inequality only gives the weaker statement

$$P(|X - \mu| \geq 2\sigma) \leq \frac{\sigma^2}{(2\sigma)^2} = 25\%$$

Solution of Problem 24

The expected value of X is $\mu = np = 5$. The standard deviation of X is

$$\sigma = \sqrt{npq} = \sqrt{2.5} = 1.6$$

The probability that X takes a value more than two standard deviations from μ is

$$P(|X - \mu| \geq 4) = 2.1\%$$

.....Alcatel-Lucent 

www.alcatel-lucent.com/careers

What if you could build your future and create the future?

One generation's transformation is the next's status quo. In the near future, people may soon think it's strange that devices ever had to be "plugged in." To obtain that status, there needs to be "The Shift".



In contrast, Chebyshev’s Inequality only gives the weaker statement

$$P(|X - \mu| \geq 2\sigma) \leq \frac{\sigma^2}{(2\sigma)^2} = 25\%$$

Solution of Problem 25

The expected value of X is $\mu = \lambda$ and the standard deviation is $\sigma = \sqrt{\lambda} = 3.2$. The probability that X takes a value more than two standard deviations from μ is

$$P(X < 4) + P(X > 16) = 3.7\%$$

In contrast, Chebyshev’s Inequality only gives the weaker statement

$$P(|X - \mu| \geq 2\sigma) \leq \frac{\sigma^2}{(2\sigma)^2} = 25\%$$

Solution of Problem 26

The expected value of X is $q/p = 1$ and the standard deviation is $\sigma = \sqrt{q/p^2} = 1.4$. The probability that X takes a value more than two standard deviations from μ is

$$P(X \geq 4) = \left(\frac{1}{2}\right)^4 = 6.3\%$$

In contrast, Chebyshev’s Inequality only gives the weaker statement

$$P(|X - \mu| \geq 2\sigma) \leq \frac{\sigma^2}{(2\sigma)^2} = 25\%$$

Solution of Problem 27

The observations are ordered according to size:

$$2, 2, 4, 5, 7, 9, 9, 10, 12, 20$$

The median is the mean of the two “middle” observations, i.e.

$$x(0.5) = \frac{7 + 9}{2} = 8$$

Similarly the lower and upper quartiles are

$$x(0.25) = \frac{2 + 4}{2} = 3, \quad x(0.75) = \frac{10 + 12}{2} = 11$$

The inter-quartile range thus becomes $11 - 3 = 8$.

Solution of Problem 28

The mean is

$$\bar{x} = \frac{2 + 5 + 10 + 11}{4} = 7$$

The empirical variance is

$$s^2 = \frac{(2 - 7)^2 + (5 - 7)^2 + (10 - 7)^2 + (11 - 7)^2}{4 - 1} = 18$$

The empirical standard deviation thus becomes

$$s = \sqrt{18} \approx 4.24$$

Solution of Problem 29

If the coin is fair, then k originates from a $\text{Bin}(10, \frac{1}{2})$ distribution. A $\text{Bin}(10, \frac{1}{2})$ distributed random variable X has the following point probabilities:

k	...	5	6	7	8	9	10
$P(X = k)$...	24.6%	20.5%	11.7%	4.4%	1.0%	0.1%

The test's significance probability P is the probability of having k or more “heads” when tossing a fair coin ten times, i.e.

$$P = P(X \geq k)$$

For $k = 8$ we get $P = 5.5\%$ and \mathbf{H}_0 must be accepted. For $k = 9$ we get $P = 1.1\%$. A test at significance level 5% should therefore reject \mathbf{H}_0 if $k \geq 9$, and accept \mathbf{H}_0 if $k \leq 8$. The risk of committing an error of type I is thus 1.1%.

If the true value of p is 0.75, then the strength of the test is

$$P(Y \geq 9) = 24.4\%$$

and the risk of an error of type II is

$$P(Y \leq 8) = 75.6\%$$

Here Y is a $\text{Bin}(10; 0.75)$ distributed random variable.

Solution of Problem 30

The number X of heads is obviously $\text{Bin}(10, \frac{1}{2})$ distributed. In order to calculate P_1 we simply use the point probabilities since this is doable in this case, and moreover we are at the borderline with respect to when the normal approximation may be used. We get

$$\begin{aligned} P_1 &= P(X = 6) + P(X = 7) + P(X = 8) + P(X = 9) + P(X = 10) \\ &= 0.205 + 0.117 + 0.044 + 0.010 + 0.001 \\ &= 37.7\% \end{aligned}$$

Solution of Problem 31

The number X of heads is obviously $\text{Bin}(100, \frac{1}{2})$ distributed. It is now advantageous to use the normal approximation to the binomial distribution, and we get

$$P_2 = P(X \geq 60) \approx 1 - \Phi\left(\frac{9.5}{\sqrt{25}}\right) = 1 - \Phi(1.9)$$

Table C.2 shows

$$P_2 = 1 - 0.971 = 2.9\%$$

Solution of Problem 32

The number X of heads is $\text{Bin}(1000, \frac{1}{2})$ distributed. We use the normal approximation again:

$$P_3 = P(X \geq 600) \approx 1 - \Phi\left(\frac{99.5}{\sqrt{250}}\right) = 1 - \Phi(6.29)$$

Table C.2 shows that P_3 is (much) less than 0.01%.

Solution of Problem 33

The number X of customers during an hour may be assumed to be Poisson distributed with intensity $\lambda = 10$, i.e. $X \sim \text{Pois}(10)$. The sought-after probability

$$P = P(X \leq 8) = \sum_{i=0}^8 P(X = i)$$

is calculated using point probabilities of the Poisson distribution (section 9.3):

$$P = 33.2\%$$



Maastricht University

Leading in Learning!

**Join the best at
the Maastricht University
School of Business and
Economics!**

Top master's programmes

- 33rd place Financial Times worldwide ranking: MSc International Business
- 1st place: MSc International Business
- 1st place: MSc Financial Economics
- 2nd place: MSc Management of Learning
- 2nd place: MSc Economics
- 2nd place: MSc Econometrics and Operations Research
- 2nd place: MSc Global Supply Chain Management and Change

Sources: Keuzegids Master ranking 2013; Elsevier 'Beste Studies' ranking 2012; Financial Times Global Masters in Management ranking 2012

Maastricht University is the best specialist university in the Netherlands (Elsevier)

**Visit us and find out why we are the best!
Master's Open Day: 22 February 2014**

www.mastersopenday.nl



Solution of Problem 34

The number Y of customers during an entire day is the sum of ten independent $\text{Pois}(10)$ distributed random variables:

$$Y = X_1 + \cdots + X_{10}$$

Thus Y is itself Poisson distributed with intensity 100, cf. section 9.5, and thus $Y \sim \text{Pois}(100)$ (one could also argue more directly to show this). Our probability

$$Q = P(Y \leq 80)$$

is calculated using the normal approximation to the Poisson distribution:

$$Q \approx \Phi\left(\frac{-19.5}{\sqrt{100}}\right) = \Phi(-1.95) = 2.6\%$$

where we looked up Φ in Table C.2.

Solution of Problem 35

If O_i denotes the number of games with i goals, then the observations can be presented in a table as follows:

i	O_i
0	8
1	13
2	18
3	11
4	10
5	2
6	2

We have to investigate whether these observations could originate from a $\text{Pois}(\lambda)$ distribution. The intensity λ is estimated as $144/64 = 2.25$ since a total of 144 goals were scored in the 64 games. The point probabilities in a $\text{Pois}(2.25)$ distribution are

i	p_i
0	0.105
1	0.237
2	0.267
3	0.200
4	0.113
5	0.051
6	0.019
≥ 7	0.008

The expected numbers thus become (since $E_i = 64p_i$)

i	E_i
0	6.7
1	15.2
2	17.1
3	12.8
4	7.2
≥ 5	4.9

Note that we have merged some categories here in order to get $E_i \geq 3$. Now we compute the statistic:

$$\begin{aligned} \chi^2 &= \frac{(8 - 6.7)^2}{6.7} + \frac{(13 - 15.2)^2}{15.2} + \frac{(18 - 17.1)^2}{17.1} + \\ &\quad \frac{(11 - 12.8)^2}{12.8} + \frac{(10 - 7.2)^2}{7.2} + \frac{(4 - 4.9)^2}{4.9} \\ &= 2.1 \end{aligned}$$

Since there are six categories and we have estimated one parameter from the data, the statistic must be compared with the χ^2 distribution with $df = 6 - 1 - 1 = 4$ degrees of freedom. Table C.3 gives a significance probability of more than 50%.

Conclusion: *It is reasonable to claim that the number of goals per game is Poisson distributed.*

Solution of Problem 36

The expected wait until success is the reciprocal probability of success:

$$E(V) = \frac{1}{1/6} = 6$$

The number of failures before success, $W = V - 1$, is $\text{Geo}(1/6)$ distributed. W and V have the same variance:

$$\text{var}(V) = \text{var}(W) = q/p^2 = 30$$

Solution of Problem 37

We get

$$P(W < n) = 1 - P(W \geq n) = 1 - q^n$$

where as always $q = 1 - p$.

Solution of Problem 38

The significance probability, i.e. the probability of having to use at least n throws given \mathbf{H}_0 , is

$$P = \left(\frac{5}{6}\right)^{n-1}$$

P is less than 5% for $n \geq 18$. Therefore, we have to reject \mathbf{H}_0 if n is at least 18.

Solution of Problem 39

The number Y of even numbers is hypergeometrically distributed with parameters $n = 7, r = 19, s = 18$ and $N = 37$, i.e. $Y \sim \text{HG}(7, 19, 37)$. The probability P is computed using the formula for the point probabilities:

$$P = P(Y = k) = \frac{\binom{19}{k} \cdot \binom{18}{7-k}}{\binom{37}{7}}$$

We get (in percentages):

k	0	1	2	3	4	5	6	7
$P(Y = k)$	0.3	3.4	14.2	28.8	30.7	17.3	4.7	0.5

Solution of Problem 40

A random variable $Y \sim \text{HG}(7, 19, 37)$ has expected value

$$E(Y) = nr/N = 7 \cdot 19/37 = 3.595$$

and standard deviation

$$\begin{aligned} \sqrt{\text{var}(Y)} &= \sqrt{nrs(N - n)/(N^2(N - 1))} \\ &= \sqrt{7 \cdot 19 \cdot 18 \cdot (37 - 7)/(37^2(37 - 1))} \\ &= 1.207. \end{aligned}$$

agence cdc © Photonstop

> Apply now

**REDEFINE YOUR FUTURE
AXA GLOBAL GRADUATE
PROGRAM 2015**

redefining / standards 



The normal approximation now gives

$$P = P(Y = k) \approx \varphi\left(\frac{k - 3.595}{1.207}\right) \cdot \frac{1}{1.207}$$

where φ is the density of the standard normal distribution (section 15.4). Using the different possible values for k , we get (in percentages):

k	0	1	2	3	4	5	6	7
$P(Y = k)$	0.4	3.3	13.8	29.3	31.2	16.8	4.5	0.6

If we compare with the previous problem, we see that the normal approximation in this case is acceptable without being magnificent. The condition that “ n be small compared to both r and s ” (section 11.6), in our case “7 is small compared to both 19 and 18”, can thus be said to be just fulfilled here.

Solution of Problem 41

The formula for the point probabilities in the multinomial distribution gives

$$P = \binom{6}{0\ 0\ 0\ 2\ 2\ 2} \cdot \left(\frac{1}{6}\right)^6 \approx 0.19\%$$

Solution of Problem 42

It is most easy to compute ad hoc:

$$P = 1 \cdot \frac{5}{6} \cdot \frac{4}{6} \cdot \frac{3}{6} \cdot \frac{2}{6} \cdot \frac{1}{6} \approx 1.5\%$$

Solution of Problem 43

The average number of goals per game is

$$\bar{k} = 2.250$$

and the empirical variance is

$$s^2 = 2.254$$

The parameters of the negative binomial distribution are estimated as

$$\hat{n} = \frac{\bar{k}^2}{s^2 - \bar{k}} = 1276$$

and

$$\hat{p} = \frac{\bar{k}}{s^2} = 0.998$$

Now we can calculate the expected values:

i	E_i
0	6.8
1	15.2
2	17.1
3	12.8
4	7.2
≥ 5	5.0

These expected values are almost identical with those from Problem 35. Since the Poisson distribution is a more natural choice and also requires one parameter less, there is no reason to explain the observations using a negative binomial distribution.

Solution of Problem 44

A’s lifespan is the wait until an event (death) that occurs spontaneously with intensity $\lambda_A = 1/5$. Similarly, B’s lifespan is the wait until an event that occurs spontaneously with intensity $\lambda_B = 1/10$. The lifespan of the device is thus the wait until an event that occurs spontaneously with intensity $\lambda = \lambda_A + \lambda_B = 3/10$. The expected lifespan of the device thus becomes $\lambda^{-1} \approx 3.3$ years.

Solution of Problem 45

The expected wait until the first component dies is seen to be 2.5 years as in the previous problem. After this, the expected wait until the second component dies is seen to be five years. The expected lifespan of the device thus becomes 7.5 years.

Solution of Problem 46

The standard deviation is $\sigma = 2$. We get

$$P(X \geq 6) = \Phi\left(-\frac{6 - \mu}{\sigma}\right) = \Phi(-1.5) = 6.68\%$$

by looking up in Table C.2.

Solution of Problem 47

The information $\Phi(u) = P(X \leq 0) = 0.10$ implies $u = -1.28$ (Table C.2). This yields $0 = 5 - 1.28 \cdot \sigma$ and thereby $\sigma = 5/1.28 = 3.91$. The variance thus becomes $\sigma^2 = 15.3$.

Solution of Problem 48

The information $P(X \leq 0) = 0.40$ implies

$$\Phi\left(\frac{0 - \mu}{\sigma}\right) = 0.40 \text{ and thus } \frac{0 - \mu}{\sigma} = -0.25$$

The information $P(X \geq 10) = 0.10$ implies

$$\Phi\left(-\frac{10 - \mu}{\sigma}\right) = 0.10 \text{ and thus } \frac{10 - \mu}{\sigma} = 1.28$$

These two equations together yield $\mu = 1.6$ and $\sigma = 6.5$.

Solution of Problem 49

$X + Y$ is itself normally distributed with expected value $\mu = 1 + 2 = 3$ and variance $\sigma^2 = 3 + 4 = 7$ (section 15.10). Therefore

$$P(X + Y \leq 5) = \Phi\left(\frac{5 - 3}{\sqrt{7}}\right) = \Phi(0.76) = 78\%$$

Solution of Problem 50

The expected value is 70, and the variance 140. Table C.3 shows that $P(Q < 100)$ is a bit below 99%.

Solution of Problem 51

The sum is χ^2 distributed with $df = 10$ degrees of freedom. Table C.3 shows that the desired probability lies somewhere between 80% and 90%.

Solution of Problem 52

The expected value is 0, and the variance 2. Table C.4 shows that $P(T < 2)$ is between 90% and 95%.

Solution of Problem 53

Table C.5 shows $x = 2.88$.

Solution of Problem 54

We calculate the statistic

$$u = \frac{\sqrt{10} \cdot 0.7}{\sqrt{3}} = 1.28$$

Since we test H_0 against all possible alternative hypotheses, the significance probability becomes

$$P = 2 \cdot \Phi(-u) = 20\%$$

If we, as commonly, test at significance level 5%, then H_0 cannot be rejected.

Solution of Problem 55

The statistic is

$$u = \frac{\sqrt{n} \cdot 0.7}{\sqrt{3}}$$

If the significance probability

$$P = 2 \cdot \Phi(-u)$$

is to be less than 5%, then u must be greater than 1.96. This implies

$$n > \left(\frac{1.96 \cdot \sqrt{3}}{0.7} \right)^2 \approx 23$$

Solution of Problem 56

The obvious thing to do is to use *Student's t-test* in this situation. We have earlier (Problem 28) computed the mean $\bar{x} = 7$ and the empirical standard deviation $s = 4.24$ of the sample. The statistic now becomes

$$t = \frac{\sqrt{4}(\bar{x} - 0)}{s} = 3.30$$

The significance probability thus becomes

$$1 - F_{\text{Student}}(3.30) < 1 - 0.975 = 2.5\%$$

where F_{Student} is the distribution function for Student's t distribution with $df = 4 - 1 = 3$ degrees of freedom. It is seen in Table C.4 that $F_{\text{Student}}(3.30)$ is a bit above 0.975. Since the significance probability thereby is less than 5%, we reject \mathbf{H}_0 after testing against \mathbf{H}_1 .

Solution of Problem 57

We have earlier (Problem 28) computed the mean $\bar{x} = 7$ and the empirical variance $s^2 = 18$ of the sample. The relevant statistic (section 17.3) is

$$q = \frac{(4 - 1)s^2}{10} = 5.4$$

The significance probability thus becomes

$$1 - F_{\chi^2}(5.4) \approx 1 - 0.85 = 15\%$$

where F_{χ^2} is the distribution function of the χ^2 distribution with $df = 4 - 1 = 3$ degrees of freedom. It is seen in Table C.3 that $F_{\chi^2}(5.4)$ is more or less in the middle between 0.8 and 0.9. Since the significance probability thereby is greater than 5%, \mathbf{H}_0 cannot be rejected – even though

Empowering People. Improving Business.

BI Norwegian Business School is one of Europe's largest business schools welcoming more than 20,000 students. Our programmes provide a stimulating and multi-cultural learning environment with an international outlook ultimately providing students with professional skills to meet the increasing needs of businesses.

BI offers four different two-year, full-time Master of Science (MSc) programmes that are taught entirely in English and have been designed to provide professional skills to meet the increasing need of businesses. The MSc programmes provide a stimulating and multi-cultural learning environment to give you the best platform to launch into your career.

- MSc in Business
- MSc in Financial Economics
- MSc in Strategic Marketing Management
- MSc in Leadership and Organisational Psychology

BI NORWEGIAN BUSINESS SCHOOL

EFMD EQUIS ACCREDITED

www.bi.edu/master



the empirical variance 18 is markedly greater than 10; the explanation lies in the very small size of the sample.

Solution of Problem 58

If the variances σ_1^2 and σ_2^2 are different, then the problem is unsolvable (this is the so-called *Fisher-Behrens problem* mentioned in the Compendium). Therefore, we first have to test the auxiliary hypothesis

$$\mathbf{H}_0^* : \sigma_1 = \sigma_2$$

against the alternative

$$\mathbf{H}_1^* : \sigma_1 \neq \sigma_2$$

Only if \mathbf{H}_0^* is accepted, the original problem can be solved.

The first sample has mean $\bar{x} = 7$ and empirical variance $s_1^2 = 18$. The second sample has mean $\bar{y} = 13$ and empirical variance $s_2^2 = 15.3$. We calculate the statistic (section 17.7 and 17.9)

$$v = \frac{s_1^2}{s_2^2} = 1.17$$

as well as

$$v^* = \max \left\{ v, \frac{1}{v} \right\} = 1.17$$

The significance probability is

$$P = 2 \cdot (1 - F_{\text{Fisher}}(v^*))$$

where F_{Fisher} is the distribution function of Fisher's F distribution with $4 - 1 = 3$ degrees of freedom in both numerator and denominator. Table C.5 gives $F_{\text{Fisher}}(5.39) = 90\%$ which implies $F_{\text{Fisher}}(1.17) < 90\%$. We thus get

$$P > 2 \cdot (1 - 90\%) = 20\%$$

With a significance probability as large as that, we *accept* the auxiliary hypothesis \mathbf{H}_0^* .

Now we are ready to test \mathbf{H}_0 against \mathbf{H}_1 . The procedure is described in section 17.8. The “pooled” variance is computed as

$$s_{\text{pool}}^2 = \frac{3s_1^2 + 3s_2^2}{6} = 16.7$$

The statistic thus becomes

$$t = \frac{\bar{x} - \bar{y}}{\sqrt{(1/4 + 1/4)s_{\text{pool}}^2}} = \frac{7 - 13}{\sqrt{(1/4 + 1/4)16.7}} = -2.08$$

The significance probability is now seen to be

$$P = 1 - F_{\text{Student}}(-t) = 1 - F_{\text{Student}}(2.08) < 5\%$$

where we have looked up $F_{\text{Student}}(2.08)$ in Table C.4 under $df = 4 + 4 - 2 = 6$. Since P is less than 5%, we *reject* \mathbf{H}_0 .

Solution of Problem 59

We have $k = 3$ samples with $n_i = 4$ observations each, in total $n = 12$ observations. The means of the samples are

$$\bar{x}_1 = 7, \bar{x}_2 = 12, \bar{x}_3 = 8$$

and the empirical variances are

$$s_1^2 = 18, s_2^2 = 15.3, s_3^2 = 14$$

The grand mean is

$$\bar{x} = \frac{7 + 12 + 8}{3} = 9$$

Next we estimate the variance **within** the samples

$$s_I^2 = \frac{1}{n - k} \sum_{j=1}^3 (4 - 1)s_j^2 = \frac{18 + 15.3 + 14}{3} = 15.4$$

and the variance **between** the samples

$$s_M^2 = \frac{1}{k - 1} \sum_{j=1}^3 4(\bar{x}_j - \bar{x})^2 = 2((7 - 9)^2 + (12 - 9)^2 + (8 - 9)^2) = 28$$

Finally the statistic can be computed:

$$v = \frac{s_M^2}{s_I^2} = \frac{28}{15.4} = 1.81$$

The significance probability is

$$P = 1 - F_{\text{Fisher}}(v) = 1 - F_{\text{Fisher}}(1.81)$$

where F_{Fisher} is the distribution function of Fisher's F distribution with $k - 1 = 2$ degrees of freedom in the numerator and $n - k = 9$ degrees of freedom in the denominator. Table C.5 shows $F_{\text{Fisher}}(1.81) < 90\%$ and thus $P > 10\%$. Consequently, we accept \mathbf{H}_0 .

Let us finally sum up the calculations in the usual ANOVA table:

sample number	1	2	3
	2	7	3
	5	11	8
	10	14	9
	11	16	12
Mean \bar{x}_j	7	12	8
Empirical variance s_j^2	18	15.3	14
$\bar{x} = 9$	(grand mean)		
$s_I^2 = (s_1^2 + s_2^2 + s_3^2)/3 = 15.4$	(variance within samples)		
$s_M^2 = 2 \sum (\bar{x}_j - \bar{x})^2 = 28$	(variance between samples)		
$v = s_M^2/s_I^2 = 1.81$	(statistic)		

Solution of Problem 60

The *observed* numbers O_i are the numbers of the second table. The *expected* numbers E_i are easy to find:

	E_i
Berlingske Tidende	14
Politiken	17
Jyllands-Posten	20
Information	5
B.T.	25
Ekstra Bladet	19

The statistic thus becomes

$$\chi^2 = \frac{(18 - 14)^2}{14} + \frac{(13 - 17)^2}{17} + \frac{(22 - 20)^2}{20} + \frac{(2 - 5)^2}{5} + \frac{(16 - 25)^2}{25} + \frac{(29 - 19)^2}{19} = 12.6$$

In order to find the significance probability P of the test, we use the χ^2 distribution with $df = 6 - 1 = 5$ (the degrees of freedom are one less than the number of categories). The result is (by looking up in Table C.3):

$$P = 1 - F_{\chi^2}(12.6) \approx 4\%$$

Need help with your dissertation?

Get in-depth feedback & advice from experts in your topic area. Find out what you can do to improve the quality of your dissertation!

[Get Help Now](#)



Go to www.helpmyassignment.co.uk for more info



Since the significance probability is less than 5%, we conclude that the ratios have changed significantly.

Solution of Problem 61

We compute the *standardized residuals*:

	r_i
Berlingske Tidende	1.2
Politiken	-1.1
Jyllands-Posten	0.5
Information	-1.4
B.T.	-2.1
Ekstra Bladet	2.5

Since standardized residuals numerically greater than 2 are a sign of an extreme observed number, we conclude that *B.T.* has decreased markedly, whereas *Ekstra Bladet* has increased markedly.

Solution of Problem 62

A total of $N = 99953$ cars were observed which gives a mean of $\lambda = 14279$ cars per day. If the seven observed numbers are from the same distribution (e.g. a $\text{Pois}(14279)$ distribution), the expected numbers will be

	E_i
Monday	14279
Tuesday	14279
Wednesday	14279
Thursday	14279
Friday	14279
Saturday	14279
Sunday	14279

The statistic thus becomes

$$\chi^2 = \sum_{i=1}^7 \frac{(O_i - E_i)^2}{E_i} = 113.3$$

This must be compared with the χ^2 distribution with $df = 7 - 1 - 1 = 5$ degrees of freedom (since we have estimated one parameter λ from the observations). We get a significance probability

$$P = 1 - F_{\chi^2}(113.3)$$

far below 0.1%, and can clearly reject the hypothesis that the numbers were from the same distribution.

Solution of Problem 63

What we have is a contingency table with two rows and two columns, i.e. a 2×2 table. In total, there are $N = 223$ observations. If there is independence between rows and columns, the expected

numbers will be

	pro	contra
Danes	63.9	28.1
Swedes	91.1	39.9

since the expected number in, say, the upper left cell is

$$E_{11} = \frac{R_1 S_1}{N} = \frac{92 \cdot 155}{223} = 63.9$$

The statistic thus becomes

$$\chi^2 = \left(\frac{70 \cdot 46 - 22 \cdot 85}{223} \right)^2 \left(\frac{1}{63.9} + \frac{1}{28.1} + \frac{1}{91.1} + \frac{1}{39.9} \right) = 3.20$$

The significance probability of the test thus becomes

$$P = 1 - F_{\chi^2}(3.20) \approx 20\%$$

where we have used the χ^2 distribution with $df = 1$ degree of freedom. Since P is greater than 5%, we *cannot* detect any regional dependence in the opinion poll.

Solution of Problem 64

Brain power

By 2020, wind could provide one-tenth of our planet's electricity needs. Already today, SKF's innovative know-how is crucial to running a large proportion of the world's wind turbines.

Up to 25 % of the generating costs relate to maintenance. These can be reduced dramatically thanks to our systems for on-line condition monitoring and automatic lubrication. We help make it more economical to create cleaner, cheaper energy out of thin air.

By sharing our experience, expertise, and creativity, industries can boost performance beyond expectations. Therefore we need the best employees who can meet this challenge!

The Power of Knowledge Engineering

Plug into The Power of Knowledge Engineering. Visit us at www.skf.com/knowledge

SKF



It is reasonable to perform a one-sided test, i.e. to test the null hypothesis

$$\mathbf{H}_0 : \text{no effect of medicine}$$

against the alternative hypothesis

$$\mathbf{H}_1 : \text{positive effect of medicine}$$

Given \mathbf{H}_0 the expected numbers become

	fit	ill
medicine	5	15
placebo	5	15

We now calculate the one-sided statistic

$$u = \left(\frac{8 \cdot 18 - 2 \cdot 12}{40} \right) \sqrt{\left(\frac{1}{5} + \frac{1}{15} + \frac{1}{5} + \frac{1}{15} \right)} = 2.19$$

Given \mathbf{H}_0 , u will be standard normally distributed, whereas a positive value of u will be expected given \mathbf{H}_1 . The significance probability thus becomes

$$1 - \Phi(2.19) = 1.4\%$$

and thereby we reject \mathbf{H}_0 in favour of \mathbf{H}_1 .

Solution of Problem 65

Let us use a one-sided test, i.e. test the null hypothesis

$$\mathbf{H}_0 : \text{women watch as much football as men}$$

against the alternative hypothesis

$$\mathbf{H}_1 : \text{men watch more football than women}$$

If we present the observations in a 2×2 table, it looks like this:

	yes	no
men	3	0
women	0	3

Given \mathbf{H}_0 , the expected numbers are

	yes	no
men	1.5	1.5
women	1.5	1.5

Since these are less than 5, we *cannot* use a χ^2 -test. Instead we use Fisher’s exact test. The significance probability thus becomes

$$P_{\text{Fisher}} = \frac{3!3!3!3!}{6!3!0!0!3!} = 5\% \text{ (exact!)}$$

Therefore, we can reject H_0 at significance level 5%.

Solution of Problem 66

We have to use Wilcoxon’s test for one set of observations. First we calculate the differences $d_i = x_i - y_i$ and assign a *rank* to each of them:

Plant no.	x_i	y_i	d_i	rank
1	41	27	14	9
2	51	59	-8	5
3	66	76	-10	7.5
4	68	65	3	2
5	46	36	10	7.5
6	69	54	15	10
7	47	49	-2	1
8	44	51	-7	4
9	60	55	5	3
10	44	35	9	6

Note: since no. 3 and no. 5 have the same numerical value, they were both given their average rank.

We now calculate the two statistics:

$$t_+ = 9 + 2 + 7.5 + 10 + 3 + 6 = 37.5 \text{ and } t_- = 5 + 7.5 + 1 + 4 = 17.5$$

At this point we can check that $t_+ + t_- = 55 = 10 \cdot (10 + 1)/2$, which shows that we have computed correctly.

If the pesticide has had an effect, the d_i ’s should be mainly positive. So we have to investigate whether t_- is “extremely” small. Table C.8 shows that the significance probability is considerably more than 5%. Therefore, we cannot prove any significant effect of the pesticide.

Solution of Problem 67

We can use the normal approximation and seek the standard normally distributed statistic

$$z = \frac{t_+ - \mu}{\sigma}$$

Since

$$\mu = \frac{100 \cdot 101}{4} = 2525 \text{ and } \sigma = \sqrt{\frac{100 \cdot 101 \cdot 201}{24}} = 290.8$$

we get

$$z = 1.79$$

The significance probability thereby becomes

$$1 - \Phi(1.79) \approx 4\%$$

and we can conclude that the pesticide *has* had a significant effect.

Solution of Problem 68

We have to use Wilcoxon’s test for two sets of observations. So we have $n = 14$ observations in the first set and $m = 11$ observations in the second set. Each observation is given a rank between 1 and $n + m = 25$:

Car no.	Line 1	Line 2
1	18	7
2	23	11
3	3	22
4	15	13
5	9	17
6	2	21
7	20	4
8	1	16
9	10	8
10	19	24
11	6	25
12	14	
13	5	
14	12	

What do you want to do?

No matter what you want out of your future career, an employer with a broad range of operations in a load of countries will always be the ticket. Working within the Volvo Group means more than 100,000 friends and colleagues in more than 185 countries all over the world. We offer graduates great career opportunities – check out the Career section at our web site www.volvogroup.com. We look forward to getting to know you!

VOLVO
 AB Volvo (publ)
www.volvogroup.com

VOLVO TRUCKS | RENAULT TRUCKS | MACK TRUCKS | VOLVO BUSES | VOLVO CONSTRUCTION EQUIPMENT | VOLVO PENTA | VOLVO AERO | VOLVO IT
 VOLVO FINANCIAL SERVICES | VOLVO 3P | VOLVO POWERTRAIN | VOLVO PARTS | VOLVO TECHNOLOGY | VOLVO LOGISTICS | BUSINESS AREA ASIA



The statistic t_x is the sum of the n ranks corresponding to Line 1:

$$t_x = 18 + 23 + 3 + 15 + 9 + 2 + 20 + 1 + 10 + 19 + 6 + 14 + 5 + 12 = 157$$

Since we are not testing against any particular alternative hypothesis, we have to consider the minimum

$$t := \min\{t_x, n(n + m + 1) - t_x\} = \min\{157, 207\} = 157$$

We look up in Table C.9 under $n = 14$ and $m = 11$ and find the number 151. Since t is *not* less than the table value, we *cannot* prove any significant difference between the number of defects at significance level $\alpha = 10\%$.

Solution of Problem 69

We may use the normal approximation to Wilcoxon's test for two sets of observations. We know that t_x is normally distributed with expected value

$$\mu = \frac{n(n + m + 1)}{2} = 7550$$

and standard deviation

$$\sigma = \sqrt{\frac{nm(n + m + 1)}{12}} = \sqrt{62917} = 251$$

So we have to look at the standard normally distributed statistic

$$z = \frac{t_x - \mu}{\sigma} = -2.18$$

This gives a significance probability of

$$\Phi(-2.18) = 1.5\%$$

We see that there were significantly fewer defects in Line 1.